Astronomy & Astrophysics manuscript no. pap˙axj1818˙rev
August 2, 2012

© ESO 2012

arXiv:1208.0249v1 [astro-ph.HE] 1 Aug 2012

The magnetar candidate AX J1818.8−1559

S. Mereghetti[1], P. Esposito[1], A. Tiengo[2,1], D. Götz[3], G.L. Israel[4], A. De Luca[1]

[1] INAF – Istituto di Astrofisica Spaziale e Fisica Cosmica Milano, via E. Bassini 15, I-20133 Milano, Italy
[2] IUSS – Istituto Universitario di Studi Superiori, viale Lungo Ticino Sforza 56, I-27100 Pavia, Italy
[3] AIM – CEA/Irfu/Service d'Astrophysique, Orme des Merisiers, F-91191 Gif-sur-Yvette, France
[4] INAF – Osservatorio Astronomico di Roma, via Frascati 33, I-00040 Monteporzio Catone, Italy

Received June 27, 2012 / Accepted August 1, 2012

ABSTRACT

In October 2007 a hard X-ray burst was detected by the $INTEGRAL$ satellite from a direction consistent with the position of AX J1818.8−1559, an X-ray source at low Galactic latitude discovered with the *ASCA* satellite in 1996-1999. The short duration (0.8 s) and soft spectrum (power law photon index of 3.0±0.2) of the burst in the 20–100 keV range are typical of Soft Gamma-ray Repeaters and Anomalous X–ray Pulsars. We report on the results of an observation of AX J1818.8−1559 obtained with the *Suzaku* satellite in October 2011. The source spectrum, a power law with photon index ∼1.5, and flux ($\sim 2\times10^{-12}$ erg cm^{-2} s^{-1}, 2-10 keV) do not show significant variations with respect to the values derived from archival data of various satellites ($ROSAT$, *XMM–Newton*, $Chandra$, $Swift$) obtained from 1993 to 2011. We discuss possible interpretations for AX J1818.8−1559 and, based on its association with the *INTEGRAL* burst, we propose it as a new member of the small class of magnetar candidates.

Key words. X-rays: individual: AX J1818.8−1559 – Gamma-ray Burst: individual: GRB071017 – Stars: magnetars

1. Introduction

Although early suggested (e.g. Pacini 1967), the relevance of magnetic energy in powering the emission from neutron stars has been recognized only recently, with the discovery of anomalous X-ray pulsars (AXPs) and soft gamma-ray repeaters (SGRs). These are spinning-down, isolated neutron stars with luminosity in the soft and hard X-rays typically larger than their rotational energy loss and characterized by the emission of powerful bursts and flares. AXPs and SGRs were historically divided in two classes, but many observations indicate that there are no substantial differences betwen them (see Mereghetti 2008 for a review). Even though alternative interpretations have been proposed (Chatterjee et al. 2000; Alpar 2001; Cea 2006; Xu 2007; Horvath 2007; Ouyed et al. 2007; Ertan et al. 2009), the model involving magnetars, i.e. highly magnetized neutron stars, is the one that currently better explains the properties of AXPs and SGRs (Duncan & Thompson 1992; Thompson & Duncan 1995; Thompson et al. 2002). According to the magnetar model, the emission from these sources is ultimately powered by the energy stored in their strong magnetic fields, with $B \sim 10^{13}$–10^{14} G in the magnetosphere, and possibly reaching even larger values in the neutron star interior.

The originally small sample of AXPs/SGRs has increased significantly in the latest years, mainly thanks to the discovery of transient sources, which remained undetected (or unnoticed) until they underwent bright outbursts, often accompanied by the emission of short bursts (Rea & Esposito 2011). In a few other cases, new magnetar candidates were found by thorough investigation of persistent X-ray sources with peculiar properties and/or possible associations with supernova remnants (e.g. Gelfand & Gaensler 2007; Halpern & Gotthelf 2010b,a), or by the discovery of a peculiar radio pulsar (Levin et al. 2010; Anderson et al. 2012).

Many of the new additions to the magnetar family have shown that these objects present a variety of properties unsuspected from the original members of the class and which connect them with other classes of isolated neutron stars. The radio detections of XTE J1810–197 and 1E 1547–54 (Camilo et al. 2006, 2007) and the very small spin-down rate of SGR 0418+5729 (Rea et al. 2010), implying a dipole field close to that of normal pulsars, are just a few examples.

The X-ray source AX J1818.8−1559 was discovered during a survey of the Galactic plane carried out with *ASCA* in 1996–1999 (Sugizaki et al. 2001), but it did not attract particular attention until *INTEGRAL* detected a short burst from its direction in October 2007. Here we report on a detailed re-analysis of the *INTEGRAL* burst and on the results of a long X-ray observation of AX J1818.8−1559 carried out with the *Suzaku* satellite in October 2011. We also present archival X-ray data, that were mostly unpublished, to provide an exhaustive description of the X-ray properties of AX J1818.8−1559, and discuss its possible inclusion in the class of magnetar candidates.

2. The burst of 2007 October 17

A short burst at low Galactic latitude (b=–0.26), was discovered and localized in real time with the *INTEGRAL* Burst Alert System (IBAS; Mereghetti et al. 2003) on 2007 October 17. The burst, reported as GRB 071017 (Mereghetti et al. 2007), was found in the data of the *ISGRI* detector (Lebrun et al. 2003), which is the low-energy imager of the *IBIS* coded-mask instrument on board *INTEGRAL* (Ubertini et al. 2003). The burst light curves in different energy ranges (Fig. 1) show that, unlike the typical gamma-ray bursts (GRBs) observed by IBIS (Vianello et al. 2009), this event was not detected above 100 keV. The 15–40 keV light curve is characterized by a single pulse lasting ∼0.8 s, with a fast rise time starting at 00:58:08.55 UT followed by a slower decay.

Send offprint requests to: S. Mereghetti, sandro@iasf-milano.inaf.it

Table 1. Log of the observations.

Satellite	Obs.Id	date	Duration (ks)	Notes
ROSAT	rp50311n00	1993 September 12	6.7	PSPC
XMM-Newton	0152834501	2003 March 28	10.0	EPIC
Swift	00020057001	2007 October 19	10.4	XRT
Chandra	7617	2007 October 21-22	5.1	ACIS-S TE
Chandra	7616	2007 October 22	39.9	ACIS-S CC
Chandra	9609	2008 February 19	1.6	ACIS
Chandra	9610	2008 September 25	1.2	HRC-I, 12′ off-axis
Suzaku	406074010	2011 October 15-18	80	XIS
Swift	00044118001	2012 March 4	0.5	XRT

We performed a spectral analysis by extracting the ISGRI data of the first 0.2 s, corresponding to the brightest part of the burst. The results confirm that, as suggested by the light curves in different energyt bands, the burst was rather soft. Its spectrum is equally well fit by a power law with photon index 3.0 ± 0.2 or by a thermal bremsstrahlung with temperature kT=25±3 keV (Fig. 2). In both cases the average flux in the 20-100 keV energy range is $\sim 2 \times 10^{-7}$ erg cm^{-2} s^{-1}.

The burst localization that was obtained in real time by the automatic IBAS software had an error of 2.9′ and was consistent with the position of AX J1818.8−1559. We reanalyzed the ISGRI data using the most recent versions of the satellite attitude information and imaging software. Using the data in the 15-50 keV, where the signal to noise is maximum, we obtained a 11.5σ detection of the burst at the coordinates R.A.=274.731, Dec.=-15.993 (J2000), with an uncertainty of 1.8′ (90% confidence level radius). The Corresponding galactic coordinates are l=15.04, b=–0.26. The *INTEGRAL* error region is indicated by the circle in Fig. 3.

3. *XMM–Newton* archival data

The region of sky in the direction of GRB 071017 was observed for about 10 ks on 2003 March 28, during a survey of the Galactic plane carried out with the *XMM–Newton* satellite. Using the Science Analysis Software (SAS V11) we reprocessed the data of the EPIC instrument (0.2–12 keV), which consists of one pn (Turner et al. 2001) and two MOS (Strüder et al. 2001) cameras. A few X-ray sources are detected in the EPIC images, but only one, at an off-axis angle of 4′, lies in the error circle of the *INTEGRAL* burst (Fig. 3). Its coordinates are $\mathrm{R.A.} = 18^{\mathrm{h}}18^{\mathrm{m}}51\overset{\mathrm{s}}{.}5$, $\mathrm{Dec.} = -15°59′23″$ (J2000), with an uncertainty of 2″. Based on the positional coincidence, as well as on the consistent spectral properties (see next paragraph), we identify the *XMM–Newton* source with AX J1818.8−1559 (Sugizaki et al. 2001).

No other sources are detected with *XMM–Newton* inside the error circle of the burst, down to a flux limit of a few 10^{-14} erg cm^{-2} s^{-1}. A few other sources are visible in the EPIC image. A faint source is located 1′ from the southern border of the error circle, at $\mathrm{R.A.} = 18^{\mathrm{h}}18^{\mathrm{m}}57\overset{\mathrm{s}}{.}9$, Dec.=–16° 02′ 08″ (J2000). Its flux is about a factor 20 smaller than that of AX J1818.8−1559. A brighter source, at $\mathrm{R.A.} = 18^{\mathrm{h}}19^{\mathrm{m}}13\overset{\mathrm{s}}{.}6$, Dec.=–16° 01′ 20″ (J2000), can be identified with the *ASCA* source AX J181915–1527 (Sugizaki et al. 2001).

To study the spectrum of AX J1818.8−1559, we extracted the pn and MOS source counts from a circle of 40″ radius and the background spectra from source free regions of the same CCD as the source. After rebinning to have at least 30 counts in each spectral bin, we fitted simultaneously the pn and two MOS spectra. A good fit was obtained with a power law with photon index $\Gamma = 1.2 \pm 0.2$, absorption $N_{\mathrm{H}} = (3.6 \pm 0.5) \times 10^{22}$ cm^{-2}, and absorbed[1] flux 1.4×10^{-12} erg cm^{-2} s^{-1} in the 2–10 keV energy range. These results are consistent with those we obtained in a quick look analysis of the same data performed soon after the *INTEGRAL* GRB detection (Tiengo et al. 2007). The source distance is unknown, but considering its location in the direction of the Galactic center region and the large absorption, we assume in the following a reference value of 10 kpc. The 0.5–10 keV luminosity, corrected for the absorption, is $\sim$2.4 $\times$ 10^{34} erg s^{-1}. An equally good fit is obtained with a blackbody of temperature $kT_{BB} = 1.9 \pm 0.1$ keV and emission radius $R_{\mathrm{BB}} = 0.12 \pm 0.05$ km. In this case the absorption is $N_{\mathrm{H}} = (1.6 \pm 0.3) \times 10^{22}$ cm^{-2}. For comparison, the total Galactic absorption in this direction is 1.2$\times$$10^{22}$ cm^{-2} (Kalberla et al. 2005). All the spectral results are summarized in Table 2. The values obtained with the power law model are very similar to those measured with *ASCA* in 1996–1999 (Sugizaki et al. 2001).

We searched for pulsations in the EPIC data with Fast Fourier Transform (FFT) and Rayleigh test methods without finding any statistically significant signal. However, this result is not particularly constraining due to the relatively small number of counts. The 3σ upper limits on the pulsed fraction, A, in the 1-10 keV energy band are of 50% for periods in the 0.3-1000 s range and of $\sim$50-70% between 145 and 300 ms[2]..

4. *Suzaku* observation

We performed a *Suzaku* observation of AX J1818.8−1559 from 2011 October 15 at 13:17 UT to October 18 at 03:32 UT. This provided about 80 ks of exposure in the XIS instrument. The XIS consists of four telescopes with a spatial resolution of about 2′ coupled to four CCD cameras operating in the 0.2–12 keV energy range (Koyama et al. 2007). At the time of our observation only the two front illuminated XIS0 and XIS3, and the back-illuminated XIS1 were operating. The three detectors were used in 1/8 window mode, which gives a time resolution of 1 s at the expense of a reduced field of view.

For the spectral analysis we screened the data using standard criteria. In particular, we considered only events with GRADE = 0,2–4,6 and excluded time intervals within ΔT = 436 s from the South Atlantic Anomaly (SAA) and pointed at elevation angles

---

[1] We use the PHABS absorption model in XSPEC with abundances from Anders & Grevesse (1989).

[2] Here and in the following we assume a sinusoidal modulation of the flux, F(t)=F$_o$ [1+Asin(Ωt)] and define the pulsed fraction as the sinusoid amplitude A. All the upper limits on A are computed as described in Israel & Stella (1996)



Fig. 1. Light curves of the 2007 October 17 burst measured with ISGRI in three energy ranges. The bin size is 50 ms.



Fig. 2. ISGRI spectrum of the 2007 October 17 burst. Top panel: data and best fit power law model. Bottom panel: residuals in units of sigma.

smaller than 5° or 20° above the night or day Earth rim, respectively. Source and background spectra were extracted from circles of 1.1′ radius and rebinned so as to have a minimum of 20 counts per spectral bin. Slightly less stringent criteria were used for the timing analysis (ΔT = 60 s, 10° day Earth minimum elevation) in order to increase the exposure time, still keeping a reasonably low background.

We obtained a single spectrum by merging the XIS 0 and 3 (the two front-illuminated CCDs) data and fitted it simultaneously with the XIS1 spectrum. All the fits were done in the 1–10 keV range. Both a power law and a blackbody model gave acceptable results, with the best fit parameters reported in Table 2. These values are similar to those derived in the *XMM–Newton* observation.

We searched for periodicities in the 2–1000 s range by an FFT analysis on the merged event lists from the three instruments. No periodicities were found, even trying different energy selections. The 3σ upper limit on the 1-10 keV pulsed fraction is ∼30-40% between 2 and 40 s, and ∼25-30% above 40 s.



Fig. 3. Image of the region of AX J1818.8−1559 in the 2-10 keV energy range obtained with the *XMM–Newton* EPIC instrument. The circle (radius 1.8′) indicates the error region of the *INTEGRAL* burst.



Fig. 4. Near infrared (K-band) image of the region of AX J1818.8−1559 from the UKIDSS. The circle with radius 0.6 arcsec indicates the source position derived with *Chandra*.

5. Other archival data

5.1. *Swift*

A short observation (2 ks) with Swift/XRT was performed on 2007 October 19, about 2 days after the *INTEGRAL* detection of the burst (Evans et al. 2007). A source with count rate 0.012 ± 0.002 counts s^{-1} was detected at the coordinates $\mathrm{R.A.} =$

Table 2. Spectral results.

	XMM–Newton 2003	*Chandra* 2007	*Suzaku* 2011
Γ	1.17±0.17	1.56 ± 0.22	1.53 ± 0.14
N_H (10^{22} cm^{-2})	3.6±0.5	$4.5^{+0.6}_{-0.5}$	$6.3^{+0.8}_{-0.7}$
F^a (10^{-12} erg cm^{-2} s^{-1})	1.37±0.07	$1.02^{+0.07}_{-0.06}$	1.04 ± 0.03
F^b (10^{-12} erg cm^{-2} s^{-1})	$2.11^{+0.17}_{-0.13}$	$2.0^{+0.3}_{-0.2}$	$2.19^{+0.16}_{-0.19}$
χ^2/dof, dof	0.88, 79	1.10, 91	1.07, 155
kT_{BB} (keV)	1.87±0.12	$1.42^{+0.11}_{-0.09}$	$1.84^{+0.09}_{-0.08}$
N_H (10^{22} cm^{-2})	1.6±0.3	2.5 ± 0.3	$2.5^{+0.5}_{-0.4}$
R^c_{BB} (km)	0.12±0.05	0.16 ± 0.02	0.11 ± 0.01
F^a (10^{-12} erg cm^{-2} s^{-1})	1.26±0.07	$8.4^{+0.6}_{-0.5}$	1.00 ± 0.03
F^b (10^{-12} erg cm^{-2} s^{-1})	1.45±0.06	1.09 ± 0.05	1.21 ± 0.04
χ^2/dof, dof	0.92, 79	1.12, 91	1.07, 155

Quoted errors are at 1σ confidence level.
[a] 2–10 keV flux.
[b] 0.5–10 keV flux corrected for absorption.
[d] For $d = 10$ kpc.

$18^{\mathrm{h}}18^{\mathrm{m}}51^{\mathrm{s}}.49$, Dec. $= -15°59'20''.07$ (J2000; $4''.1$ error radius, 90% confidence) consistent with those of AX J1818.8−1559. The small number of counts is not sufficient to perform a spectral analysis. Assuming the parameters of the *XMM–Newton* power-law fit, we derived a 2–10 keV observed flux of $\sim 1.5 \times 10^{-12}$ erg cm^{-2} s^{-1}.

This region of sky was observed again with the Swift/XRT on 2012 March 04, but no source was detected within or close to the error region of the burst. The short exposure time (0.5 ks) and off-axis position result in a rather shallow upper limit on the count rate of 0.027 counts s^{-1} (3 σ c.l.), which is a factor ~ 2 above the flux level measured in October 2007.

5.2. Chandra

The *Chandra* observations of the field of AX J1818.8−1559 are listed in Table 1[3]. The 2007 observations were done with the ACIS-S instrument as a Target of Opportunity 5 days after the *INTEGRAL* burst detection. They consist of 5 ks of imaging data, with time resolution of 3.2 s, and a longer exposure in continuous clocking (CC) mode, which provides a higher time resolution (2.85 ms), but imaging only along a single direction. AX J1818.8−1559 is detected with a significance above 10σ in the imaging data, at coordinates R.A. $= 18^{\mathrm{h}}18^{\mathrm{m}}51^{\mathrm{s}}.38$ ($\pm 0^{\mathrm{s}}.06$), Dec. $= -15°59'22''.62$ ($\pm 0''.05$) (J2000); the quoted 1σ uncertainties are statistical only, and much smaller than the *Chandra* absolute astrometric accuracy, $0''.6$ (radial, 90% c.l.). We show in Fig. 4 the *Chandra* error circle overlaid on the near infrared (K-band) image from the Galactic Plane Survey, collected as a part of the United Kingdom Infrared Deep Sky Survey (UKIDSS, Lucas et al. (2008)). No counterparts are visible down to the limit K$\sim$17–18.

We extracted the spectra from the 2007 observations as follows. For the CC data, the source photons were accumulated within a 5×5 pixels region centred on AX J1818.8−1559 (one ACIS-S pixel corresponds to 0.492 arcsec); the background events were extracted from two symmetric (with respect to the source) regions along the image strip. For the imaging data, the source counts were selected in a $\sim$5-px radius circular region and the background in an annulus with radii of 20 and 30 pixels. After checking that they gave consistent results, the two spectra were fitted together yielding the best fit parameters reported in Table 2.

AX J1818.8−1559 was also observed two further times in 2008. In the first data set (2008 February) the source is detected at a significance level of 5σ, but the small statistics do not allow us to carry out a detailed analysis. We just note that the 0.3–7 keV count rate, 0.026 ± 0.004 counts s^{-1}, is in line with that observed in the 2007 ACIS-S imaging observation, 0.024 ± 0.002 counts s^{-1}. In the second data set (2008 September), a short HRC-I exposure, the source is not detected. The 3σ upper limit on the count rate is 0.026 counts s^{-1} which, adopting the *Chandra* power-law model of Table 2, translates into an upper limit on the observed 2-10 keV flux of $\sim 5 \times 10^{-12}$ erg cm^{-2} s^{-1}.

The most sensitive data for timing analysis are provided by the 2007 data in CC mode, which yielded $\sim$1400 source photons. By an FFT analysis of these data, we could set a 3σ upper limit of $\sim$40% on the 1-10 keV pulsed fraction for periods in the range 0.01–1000 s, and $\sim$40–60% for periods between 5 and 10 ms.

5.3. ROSAT

The position of AX J1818.8−1559 was serendipitously covered by a ROSAT observation on 11-12 September 1993, falling at an off-axis angle of $\sim 35'$ in the PSPC instrument. From these data, providing an exposure of $\sim$7 ks we derived a 3σ upper limit of $\sim$0.01 counts s^{-1} on the source count rate in the 0.1-2.4 keV energy range. Assuming the same spectral shape of the *XMM–Newton* observation, this corresponds to an unabsorbed flux of the order of 2×10^{-11} erg cm^{-2} s^{-1} in the 0.5-10 keV range.

6. Discussion

A subclass of GRBs, the so called X-ray Flashes (Heise et al. 2001), can have spectra as soft as observed in GRB 071017. However, in the class of short GRBs with duration less than 2 s, such soft spectra are never observed. On the other hand, the hard X-ray (E>20 keV) spectra of short bursts from SGRs are usually well described by thermal bremsstrahlung models with temperature ∼20-40 keV (Aptekar et al. 2001; Götz et al. 2004), as observed in GRB 071017. Assuming a distance of 10 kpc, the peak luminosity of the *INTEGRAL* burst is of the order of $\sim 5 \times 10^{39}$ erg s^{-1}, which is a reasonable value for a Galactic SGR. Thus, although we cannot completely exclude that GRB 071017 was a GRB seen through the Galaxy, we believe that, in view of its location in the Galactic plane and spectral properties, an explanation in terms of a Galactic AXP/SGR is much more likely.

---

[3] Another observation was performed in August 2009 (obs. ID 10503), but the position of AX J1818.8−1559 fell in the gap between the ACIS-S CCDs

The chance probability of finding an X-ray source of a given brightness inside the 10 arcmin^2 *INTEGRAL* error circle can be estimated using the LogN-LogS relations derived by Sugizaki et al. (2001) from the survey of the Galactic plane in which AX J1818.8−1559 was discovered. Including the small contribution from extragalactic sources seen through the Galactic plane, these relations give a surface density of 1.2 deg^{-2}, for sources with flux equal or greater than that of AX J1818.8−1559. This leads to a probability of $\sim 3 \times 10^{-3}$ to find AX J1818.8−1559 in the *INTEGRAL* error region by chance, which indicates its very likely association with the burst.

The X-ray spectrum of AX J1818.8−1559 is harder than that of typical SGRs and AXPs, which, when fitted with a single power law, give photon index in the range $\sim 2 - 4$ (Mereghetti et al. 2002; Kaspi & Boydstun 2010). Harder spectra have been observed in transient magnetars during the initial phases of their outbursts, but our analysis of X-ray observations of AX J1818.8−1559 obtained from 2003 to 2012, as well as the earlier *ASCA* observations (Sugizaki et al. 2001), do not show evidence for a transient behavior. Indeed all the measured fluxes, and the upper limits, indicate a steady luminosity of 2×10^{34} (d/[10 kpc])^2 erg s^{-1}, consistent with that observed in persistent magnetars. The lack of detection of pulsations is not particularly constraining, considering that many of the AXPs and SGRs have pulsed fractions smaller than the upper limits we could set on AX J1818.8−1559 (Mereghetti et al. 2002).

Only a single burst has been observed from the direction of AX J1818.8−1559, despite this region of sky in the Galactic bulge having been repeatedly observed by high-energy satellites. This contrasts with the abundance of bursts, often grouped in time intervals of spasmodic activity, observed in the most active members of the SGR family (e.g., SGR 1806–20 (Göğüş et al. 2000), SGR 1900+14 (Israel et al. 2008), SGR 1627–41 (Esposito et al. 2008), or 1E 1547.0–5408 (Mereghetti et al. 2009)). The behavior of AX J1818.8−1559 is instead similar to that of a few magnetars from which bursts were detected only sporadically, such as for example 1E 1048.1–5937 (Gavriil et al. 2002), SGR 0418+5729 (van der Horst et al. 2010), SGR 1833–0832 (Göğüş et al. 2010), or 4U 0142+61 (Gavriil et al. 2011).

We finally note that the available observations do not favour alternative explanations for the nature of AX J1818.8−1559. Coronal emission from a star can be ruled out by the lack of a bright optical counterpart, the hard spectrum, and the large absorption, inconsistent with the small distance that would be required in this hypothesis. The X-ray to optical flux ratio excludes the possibility of a high mass X-ray binary. A low mass X-ray binary or an AGN cannot be excluded, but they are not supported by the hard spectrum and lack of long term variability, respectively.

7. Conclusion

A new observation of AX J1818.8−1559 obtained with the Suzaku satellite, as well as a comprehensive analysis of archival data from different missions could not establish the real nature of this source, which is most likely associated with a SGR-like burst observed with *INTEGRAL* in October 2007. The X-ray properties of AX J1818.8−1559 are consistent with those observed in AXPs and SGRs, and we propose it as a likely new member of this class of sources deserving further study.

Acknowledgements. This work was partially supported with contributions from the agreements ASI-INAF I/009/10/0 and I/032/10/0. This research has made use of data obtained from the *Suzaku* satellite, a collaborative mission between the space agencies of Japan (JAXA) and the USA (NASA). We also made use of data and software provided by the NASA/GFSC's HEASARC, the *Chandra* X-ray Centre and the ESA's *XMM–Newton* Science Archive. We thank the referee, V. Kaspi, for her useful comments.

References

Alpar, M. A. 2001, ApJ, 554, 1245
Anders, E. & Grevesse, N. 1989, Geochim. Cosmochim. Acta, 53, 197
Anderson, G. E., Gaensler, B. M., Slane, P. O., et al. 2012, ApJ, 751, 53
Aptekar, R. L., Frederiks, D. D., Golenetskii, S. V., et al. 2001, ApJS, 137, 227
Camilo, F., Ransom, S. M., Halpern, J. P., & Reynolds, J. 2007, ApJ, 666, L93
Camilo, F., Ransom, S. M., Halpern, J. P., et al. 2006, Nature, 442, 892
Cea, P. 2006, A&A, 450, 199
Chatterjee, P., Hernquist, L., & Narayan, R. 2000, ApJ, 534, 373
Duncan, R. C. & Thompson, C. 1992, ApJ, 392, L9
Ertan, Ü., Ekşi, K. Y., Erkut, M. H., & Alpar, M. A. 2009, ApJ, 702, 1309
Esposito, P., Israel, G. L., Zane, S., et al. 2008, MNRAS, 390, L34
Evans, P. A., Starling, R. L. C., O'Brien, P. T., et al. 2007, GRB Coordinates Network, 6942, 1
Gavriil, F. P., Dib, R., & Kaspi, V. M. 2011, ApJ, 736, 138
Gavriil, F. P., Kaspi, V. M., & Woods, P. M. 2002, Nature, 419, 142
Gelfand, J. D. & Gaensler, B. M. 2007, ApJ, 667, 1111
Götz, D., Mereghetti, S., Mirabel, I. F., & Hurley, K. 2004, A&A, 417, L45
Göğüş, E., Cusumano, G., Levan, A. J., et al. 2010, ApJ, 718, 331
Göğüş, E., Woods, P. M., Kouveliotou, C., et al. 2000, ApJ, 532, L121
Halpern, J. P. & Gotthelf, E. V. 2010a, ApJ, 725, 1384
—. 2010b, ApJ, 710, 941
Heise, J., Zand, J. I., Kippen, R. M., & Woods, P. M. 2001, in Gamma-ray Bursts in the Afterglow Era, ed. E. Costa, F. Frontera, & J. Hjorth, 16
Horvath, J. E. 2007, Ap&SS, 308, 431
Israel, G. L., Romano, P., Mangano, V., et al. 2008, ApJ, 685, 1114
Israel, G. L. & Stella, L. 1996, ApJ, 468, 369
Kalberla, P. M. W., Burton, W. B., Hartmann, D., et al. 2005, A&A, 440, 775
Kaspi, V. M. & Boydstun, K. 2010, ApJ, 710, L115
Koyama, K., Tsunemi, H., Dotani, T., et al. 2007, PASJ, 59, 23
Lebrun, F., Leray, J. P., Lavocat, P., et al. 2003, A&A, 411, L141
Levin, L., Bailes, M., Bates, S., et al. 2010, ApJ, 721, L33
Lucas, P. W., Hoare, M. G., Longmore, A., et al. 2008, MNRAS, 391, 136
Mereghetti, S. 2008, A&A Rev., 15, 225
Mereghetti, S., Chiarlone, L., Israel, G. L., & Stella, L. 2002, in Neutron Stars, Pulsars, and Supernova Remnants, ed. W. Becker, H. Lesch, & J. Trümper, 29–+
Mereghetti, S., Götz, D., Borkowski, J., Walter, R., & Pedersen, H. 2003, A&A, 411, L291
Mereghetti, S., Götz, D., Weidenspointner, G., et al. 2009, ApJ, 696, L74
Mereghetti, S., Paizis, A., Gotz, D., et al. 2007, GRB Coordinates Network, 6927, 1
Ouyed, R., Leahy, D., & Niebergal, B. 2007, A&A, 473, 357
Pacini, F. 1967, Nature, 216, 567
Rea, N. & Esposito, P. 2011, in High-Energy Emission from Pulsars and their Systems, ed. D. F. Torres & N. Rea, 247
Rea, N., Esposito, P., Turolla, R., et al. 2010, Science, 330, 944
Strüder, L., Briel, U., Dennerl, K., et al. 2001, A&A, 365, L18
Sugizaki, M., Mitsuda, K., Kaneda, H., et al. 2001, ApJS, 134, 77
Thompson, C. & Duncan, R. C. 1995, MNRAS, 275, 255
Thompson, C., Lyutikov, M., & Kulkarni, S. R. 2002, ApJ, 574, 332
Tiengo, A., Mereghetti, S., Esposito, P., De Luca, A., & Gotz, D. 2007, The Astronomer's Telegram, 1243, 1
Turner, M. J. L., Abbey, A., Arnaud, M., et al. 2001, A&A, 365, L27
Ubertini, P., Lebrun, F., Di Cocco, G., et al. 2003, A&A, 411, L131
van der Horst, A. J., Connaughton, V., Kouveliotou, C., et al. 2010, ApJ, 711, L1

Vianello, G., Götz, D., & Mereghetti, S. 2009, A&A, 495, 1005
Xu, R. 2007, Advances in Space Research, 40, 1453